



Let we have four points  $A, B, C, D \in \mathbb{R}\mathbb{P}^2$ , which are collinear, i.e. they belong to the same projective line.

Calculate their cross-ratio.

$$A = [8:20:4] \quad \text{homogeneous coord.}$$

$$u = \frac{8}{4} = 2, \quad v = \frac{20}{4} = 5 \quad \text{affine coordinates}$$

$$B = [8:18:2]$$

$$u = \frac{8}{2} = 4, \quad v = \frac{18}{2} = 9$$

$$C = [6:14:2]$$

$$u = \frac{6}{2} = 3, \quad v = \frac{14}{2} = 7$$

$$D = [x:2x+1:1]$$

$$u = \frac{x}{1} = x, \quad v = \frac{2x+1}{1} = 2x+1$$

First check that these points are really collinear.

## Lecture CXI

$$T_{ABC} = \begin{pmatrix} 8 & 8 & 6 \\ 20 & 18 & 14 \\ 4 & 2 & 2 \end{pmatrix} \sim \begin{pmatrix} 2 & 4 & 3 \\ -5 & 9 & 7 \\ 1 & 1 & 1 \end{pmatrix}$$

$$\det = 2(9-7) - 4(5-7) + 3(5-9) = 4 + 8 - 12 = 0.$$

matrix  $T_{ABC}$  is degenerate  $\iff A, B, C$  are collinear,  $C \in AB$ .

Analogously consider points  $B, C, D$ :

$$T_{BCD} = \begin{pmatrix} 8 & 6 & x \\ 18 & 14 & 2x+1 \\ 2 & 2 & 1 \end{pmatrix} \sim \begin{pmatrix} 4 & 3 & x \\ 9 & 7 & 2x+1 \\ 1 & 1 & 1 \end{pmatrix} \quad (x \neq 3)$$

$$\det = 4(7-2x-1) - 3(9-2x-1) + x(9-7) = 24 - 8x - 24 + 6x + 2x = 0.$$

Matrix  $T_{BCD}$  is degenerate  $\iff B, C, D$  are collinear,  $C \in BD$

We see that  $A, B, C$  are collinear } four points  
 $B, C, D$  are collinear }  $\Rightarrow A, B, C, D$  are collinear.

Another way to see that these points are collinear:

$$(w) - \begin{pmatrix} A \\ B \\ C \end{pmatrix} = \begin{pmatrix} 2 \\ 5 \end{pmatrix} \quad \begin{pmatrix} B \\ C \\ D \end{pmatrix} = \begin{pmatrix} 4 \\ 9 \\ x \\ 2x+1 \end{pmatrix}$$

$$w = \frac{x}{2}, V = \frac{y}{2}$$

All these points belong the line  $2w = V - 1$

$$V_A - 2w_A = 5 - 4 = V_B - 2w_B = 9 - 8 = V_C - 2w_C = 7 - 6 = V_D - 2w_D = 2x+1 - 2x = 1.$$

# Lecture C XI

To calculate the cross-ratio of collinear points  $A, B, C, D$  we choose any affine coordinate\*

$$(A, B, C, D) = \frac{(u_A - u_C)(u_B - u_D)}{(u_A - u_D)(u_B - u_C)} = \frac{(2-3)(4-x)}{(2-x)(4-3)}$$
$$(A, B, C, D) = \frac{x-4}{2-x}.$$

We can choose another coordinate

$$(A, B, C, D) = \frac{(v_A - v_C)(v_B - v_D)}{(v_A - v_D)(v_B - v_C)} = \frac{(5-7)(9-2x-1)}{(5-2x-1)(9-7)} = \frac{x-4}{2-x}.$$

We can choose  $u' = 2u + 5v \rightarrow \dots$

Cross-ratio is an invariant. Changing affine coordinate we will not change cross-ratio!

\* \* \*

Now let point  $D$  'goes' to infinity:

$$D = [x: 2x+1: 1] \xrightarrow{x \rightarrow \infty} [1; 2 + \frac{1}{x}; \frac{1}{x}] \rightarrow [1: 2: 0]$$

$D = [1: 2: 0]$  is a point at infinity which belongs to projective line  $AB$

To calculate cross-ratio:

$$(A, B, C, D) = \frac{x-4}{2-x} \rightarrow -1$$

$$(A, B, C, D) = \frac{u_A - u_C}{u_B - u_C} \text{ if } D \text{ is at infinity}$$

(See exercise 3 in the Homework 9 (C3).)

\* This coordinate has to be different for different points.

not compulsory

# Lecture CXI

## Projective transformations of $\mathbb{RP}^2$

Recall projective transformations of  $\mathbb{RP}^1$

$$K = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \quad \det K \neq 0$$

$$[x:y] \rightarrow [x':y'] = [\alpha x + \beta y : \gamma x + \delta y]$$

$$u \rightarrow u' = \frac{x'}{y'} = \frac{\alpha x + \beta y}{\gamma x + \delta y} = \frac{\alpha u + \beta}{\gamma u + \delta}$$

(See the lecture CVII)

Now define projective transform. of  $\mathbb{RP}^2$

$$K = \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix}, \det K \neq 0$$

$$[x:y:z] \rightarrow [x':y':z'] = [ax+by+cz : dx+ey+fz : gx+hy+iz]$$

$$\left( u = \frac{x}{z}, v = \frac{y}{z} \right) \rightarrow \left( u' = \frac{x'}{z'}, v' = \frac{y'}{z'} \right)$$

$$\begin{pmatrix} u \\ v \end{pmatrix} \rightarrow \begin{pmatrix} u' \\ v' \end{pmatrix} = \begin{pmatrix} \frac{x'}{z'} \\ \frac{y'}{z'} \end{pmatrix} = \begin{pmatrix} \frac{ax+by+cz}{gx+hy+iz} \\ \frac{dx+ey+fz}{gx+hy+iz} \end{pmatrix} = \begin{pmatrix} \frac{au+bv+c}{gu+hv+i} \\ \frac{du+ev+f}{gu+hv+i} \end{pmatrix}$$

Projective transformation of  $\mathbb{RP}^2$

This transformation transforms

a (projective) line to a (projective) line.

It includes usual affine transformations.

5)

## Lecture CXI

Indeed consider

$$K = \begin{pmatrix} a & b & p \\ c & d & q \\ 0 & 0 & 1 \end{pmatrix}$$

We have

$$\begin{pmatrix} u' \\ v' \end{pmatrix} = \begin{pmatrix} au + bv + p \\ cu + dv + q \\ 0 \cdot u + 0 \cdot v + 1 \end{pmatrix} = \begin{pmatrix} au + bv + p \\ cu + dv + q \\ 1 \end{pmatrix}$$

$$\begin{pmatrix} u' \\ v' \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} p \\ q \end{pmatrix}$$

This usual affine transformation.

Projective transf. transform  
 points (of  $\mathbb{RP}^2$ ) to points (of  $\mathbb{RP}^2$ )  
 likez (proj) to lines - (proj.)

Cross-ratio is invariant of projective transform.  
 Affine transformation is a transform,

of projective plane which does  
 not move projective line at  
 infinity.