

## **Introduction to Geometry (20222)**

**2019**

### **COURSEWORK**

This assignment counts for 20% of your marks.

It will take about 8 hours of work

Solutions are due by 28 March, 3pm

*Write solutions in the provided spaces.*

**STUDENT'S NAME:**

Academic Advisor (Tutor):

**1**

a) Let  $(x^1, x^2)$  be coordinates of the vector  $\mathbf{x}$ , and  $(y^1, y^2)$  be coordinates of the vector  $\mathbf{y}$  in  $\mathbf{R}^2$ .

Consider the formula

$$(\mathbf{x}, \mathbf{y}) = x^1 y^1 + x^2 y^2 + kx^1 y^2 + kx^2 y^1,$$

where  $k$  is a real parameter. Show that this formula defines a scalar product in  $\mathbf{R}^2$  in the case if  $|k| < 1$ .

Give an example of orthonormal basis for this scalar product.

Explain why this formula does not define a scalar product on  $\mathbf{R}^2$  in the case if  $|k| \geq 1$ .

[2 marks]

b) Consider the matrix  $A = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ .

Calculate the matrix  $A^9$  in the case if  $\theta = \frac{\pi}{27}$ .

Calculate the matrix  $A^{2019}$  in the case if  $\theta = \frac{\pi}{6}$ .

[1 marks]

c) In Euclidean space  $\mathbf{E}^3$  consider the following linear operator

$$A(\mathbf{x}) = \mathbf{x} + (\mathbf{a}, \mathbf{x})\mathbf{a}$$

where the vector  $\mathbf{a} = 3\mathbf{e} + 4\mathbf{f} + 12\mathbf{g}$ . Here  $\{\mathbf{e}, \mathbf{f}, \mathbf{g}\}$  is an orthonormal basis in  $\mathbf{E}^3$ .

Calculate the trace and determinant of the operator  $A$ .

[2 marks]

d) Let  $\{\mathbf{e}, \mathbf{f}\}$  be an orthonormal basis of Euclidean space  $\mathbf{E}^2$ . Consider a linear operator  $P$  such that  $\mathbf{a} = P(\mathbf{e}) = 91\mathbf{e} + 50\mathbf{f}$ ,  $\mathbf{b} = P(\mathbf{f}) = 20\mathbf{e} + 11\mathbf{f}$ .

Calculate determinant of the operator  $P$ .

Show that  $P$  is not an orthogonal operator.

Does this operator preserve an orientation of  $\mathbf{E}^2$ ? Justify your answer.

Consider the parallelogram  $\Pi_{\mathbf{a}, \mathbf{b}}$  formed by the vectors  $\mathbf{a}$  and  $\mathbf{b}$  attached at the origin.

Find the area of this parallelogram.

Show that the vertices of the parallelogram  $\Pi_{\mathbf{a},\mathbf{b}}$  are the only points of  $\Pi_{\mathbf{a},\mathbf{b}}$ , whose coordinates are both integers.

[3 marks]



**2**

We consider in this question 3-dimensional Euclidean space  $\mathbf{E}^3$ . We suppose that  $\{\mathbf{e}, \mathbf{f}, \mathbf{g}\}$  is an orthonormal basis in this space.

**a)** Let  $P$  be a linear orthogonal operator acting in  $\mathbf{E}^3$  such that its matrix in the basis  $\{\mathbf{e}, \mathbf{f}, \mathbf{g}\}$  has the following appearance

$$P = \frac{1}{7} \begin{pmatrix} 3 & * & 6 \\ -6 & -3 & 2 \\ 2 & -6 & * \end{pmatrix}.$$

Find the entries of the matrix denoted by  $*$ .

Show that the operator  $P$  preserves orientation.

We know that due to the Euler Theorem the linear operator  $P$  considered above is a rotation operator. Find the axis and the angle of this rotation.

[3 marks]

**b)** Let  $P_1$  be a rotation operator on the angle  $\theta$  around the axis directed along the vector  $\mathbf{g}$ , and  $P_2$  be a rotation operator on the same angle  $\theta$  around the axis directed along the vector  $\mathbf{e}$ :

$$\{\mathbf{e}, \mathbf{f}, \mathbf{g}\} \xrightarrow{P_1} \{\cos \theta \mathbf{e} + \sin \theta \mathbf{f}, -\sin \theta \mathbf{e} + \cos \theta \mathbf{f}, \mathbf{g}\},$$

$$\{\mathbf{e}, \mathbf{f}, \mathbf{g}\} \xrightarrow{P_2} \{\mathbf{e}, \cos \theta \mathbf{f} + \sin \theta \mathbf{g}, -\sin \theta \mathbf{f} + \cos \theta \mathbf{g}\}.$$

Show that the operator  $P = P_1 \circ P_2$  is also a rotation operator. Find the axis of rotation and the angle  $\Phi = \Phi(\theta)$  of rotation for the operator  $P$ .

Calculate the angle  $\Phi = \Phi(\theta)$  in the case  $\theta = \frac{\pi}{2}$ .

Show that in the case if  $\theta$  is small, then  $\Phi(\theta) \approx \sqrt{2}\theta$ , i.e.

$$\lim_{\theta \rightarrow 0} \frac{\Phi(\theta)}{\theta} = \sqrt{2}.$$

[5 marks]





**3**

a) Consider the curve  $\mathbf{r}(t)$ : 
$$\begin{cases} x = Rt \\ y = R\sqrt{1-t^2} \end{cases}, \quad 0 \leq t \leq 1.$$

Draw the image of this curve.

Give an example of a parameterisation of this curve with opposite orientation.

[1 marks]

b) Let  $f$  be a function in  $\mathbf{E}^2$  given by  $f = r^2 \cos 2\varphi$ , where  $r, \varphi$  are polar coordinates in  $\mathbf{E}^2$  ( $x = r \cos \varphi, y = r \sin \varphi$ ). Consider vector fields which are given in Cartesian coordinates by  $\mathbf{A} = x\partial_x + y\partial_y$ ,  $\mathbf{B} = x\partial_y - y\partial_x$ .

Calculate  $\partial_{\mathbf{A}} f$ ,  $\partial_{\mathbf{B}} f$ .

Let  $g$  be a function on  $\mathbf{E}^2$  such that differential form  $\omega = dg$  vanishes at the vector field  $\mathbf{B} = x\partial_y - y\partial_x$ :  $\omega(\mathbf{B}) \equiv 0$ . Find a function  $g$  if it is known that

$$g(x, y) \Big|_{y=0} = x^6.$$

[3 marks]