

## Homework 6

1 Calculate the integrals of the form  $\omega = xdy - ydx$  over the following three curves. Compare answers.

$$C_1: \mathbf{r}(t) \begin{cases} x = R \cos t \\ y = R \sin t \end{cases}, \quad 0 < t < \pi, \quad C_2: \mathbf{r}(t) \begin{cases} x = R \cos 4t \\ y = R \sin 4t \end{cases}, \quad 0 < t < \frac{\pi}{4}$$

$$\text{and } C_3: \mathbf{r}(t) \begin{cases} x = Rt \\ y = R\sqrt{1-t^2} \end{cases}, \quad -1 < t < 1.$$

(this exercise was done during the XIV-th lecture (Tuesday, VII-th week): see the last example in subsection 2.4 "Integration of differential forms over curves" in Lecture notes)

2 Consider an arc of parabola  $x = 2y^2 - 1$ ,  $0 < y < 1$ .

Give examples of two different parameterisations of this curve such that these parameterisations have the opposite orientation.

Calculate the integral of the form 1-form  $\omega = \sin y dx$  over this curve.

How does the answer depend on a parameterisation?

3 Calculate the integral of the form  $\omega = xdy$  over the following curves

a) closed curve  $x^2 + y^2 = 12y$

b) arc of the ellipse  $x^2 + y^2/9 = 1$  defined by the condition  $y \geq 0$ .

How does your answer depend on a choice of parameterisation?

4 a) Calculate the integrals  $\int_{C_1} \omega$  and  $\int_{C_2} \omega$  of the 1-form  $\omega = xdy - ydx$  over the curves  $C_1: x^2 + y^2 = 9$  and  $C_2: x^2 + y^2 = 6y$ .

b) Perform the calculations of integrals  $\int_{C_1} \omega$  and  $\int_{C_2} \omega$  in polar coordinates.

*Hint Performing the calculations for the curve  $C_2$  one may use the polar coordinates  $r, \varphi$  with the centre at the point  $(a, b)$ :*  $\begin{cases} x = a + r \cos \varphi \\ y = b + r \sin \varphi \end{cases}$ .

5 Calculate the integral  $\int_C \omega$  where  $\omega = xdx + ydy$  and  $C$  is

a) the straight line segment  $x = t, y = 1 - t, 0 \leq t \leq 1$

b) the segment of parabola  $x = t, y = 1 - t^n, 0 \leq t \leq 1, n = 2, 3, 4, \dots$

c) for **an arbitrary** curve starting at the point  $(0, 1)$  and ending at the point  $((1, 0))$ .

6 Show that the form 1-form  $\omega = 3x^2ydx + x^3dy$  is an exact 1-form.

Calculate integral of this form over the curves considered in exercises 2) and 3).

7. Consider in  $\mathbf{E}^2$  1-forms

a)  $xdx$ , b)  $xdy$  c)  $xdx + ydy$ , d)  $xdy + ydx$ , e)  $xdy - ydx$

f)  $x^4dy + 4x^3ydx$ .

a) Show that 1-forms a), c), d) and f) are exact forms

b) Why are 1-forms b) and e) not exact?

8 Consider 1-form

$$\omega = \frac{xdy - ydx}{x^2 + y^2}. \quad (1)$$

This form is defined in  $\mathbf{E}^2 \setminus 0$ , i.e. in all the points except origin:  $x^2 + y^2 \neq 0$ .

a) Write down this form in polar coordinates

b) † What values can take the integral  $\int_C \omega$  of this form, if  $C$  is an arbitrary curve starting at the point  $(0, 1)$  and ending at the point  $((1, 0)$  (we suppose that the curve  $C$  does not pass through the origin)

9† Let  $\omega = a(x, y)dx + b(x, y)dy$  be a closed form in  $\mathbf{E}^2$ ,  $d\omega = 0$ .

Consider the function

$$f(x, y) = x \int_0^1 a(tx, ty)dt + y \int_0^1 b(tx, ty)dt \quad (2)$$

† Show that

$$\omega = df.$$

( *This proves that an arbitrary closed form in  $\mathbf{E}^2$  is an exact form.*

† Why we cannot apply the formula (2) to the form  $\omega$  defined by the expression (1)?